arXiv:1512.05385v1 [math.FA] 11 Dec 2015

THE FRACTIONAL S-TRANSFORM ON BMO AND HARDY SPACES

BABY KALITA[1] AND SUNIL KUMAR SINGH[2,*]

Abstract. In this paper, the authors studied the fractional S-transform on BMO and Hardy spaces and generalized the results given in [15]. In introduction section, the definitions of the S-transform, fractional Fourier transform and fractional S-transform are given. In Section 2 continuity and boundedness results for the fractional S-transform in BMO and Hardy spaces are obtained. Furthermore, in Section 3 the fractional S-transform is studied on the weighted BMO_κ and weighted Hardy spaces associated with a tempered weight function.

Key words: S-transform, Fractional S-transform, BMO space, Hardy space
2010 AMS Subject Classification: 65R10; 32A37; 30H10

1. Introduction

The S-transform is a new time-frequency analysis method, which is deduced from short-time Fourier transform and an extension of wavelet transform. It was first introduced by Stockwell et al.[17] in 1996 for analyzing geophysics data and since then has been applied in several discipline, such as geophysics, oceanography, atmospheric physics, medicine, hydrogeology. The continuous S-transform of a function f with respect to the window function ω is defined as [18]

$$(S_\omega f)(\tau,\xi) = \int_{\mathbb{R}} f(t)\,\omega(\tau - t,\xi)\,e^{-i2\pi\xi t}\,dt, \text{ for } \tau,\xi \in \mathbb{R}, \tag{1.1}$$

where the window ω is assumed to satisfy the following:

$$\int_{\mathbb{R}} \omega(t,\xi)dt = 1 \text{ for all } \xi \in \mathbb{R}_0 := \mathbb{R} \setminus \{0\}. \tag{1.2}$$

The most usual window ω is a Gaussian one

$$\omega(t,\xi) = \frac{|\xi|}{k\sqrt{2\pi}}\,e^{-\xi^2 t^2/2k^2}, k > 0, \tag{1.3}$$

where ξ is the frequency, t is the time variable, and k is a scaling factor that controls the number of oscillations in the window.

Equation(1.1) can be rewritten as a convolution

$$(S_\omega f)(\tau,\xi) = \left(f(\cdot)e^{-i2\pi\xi\cdot} * \omega(\cdot,\xi)\right)(\tau). \tag{1.4}$$

Applying the convolution property for the Fourier transform in (1.4), we can obtain a direct relation between S-transform and Fourier transform as follows:

$$(S_\omega f)(\tau,\xi) = \mathscr{F}^{-1}\left\{\hat{f}(\cdot+\xi)\;\hat{\omega}(\cdot,\xi)\right\}(\tau), \tag{1.5}$$

where $\hat{f}(\eta) = (\mathscr{F}f)(\eta) = \int_{\mathbb{R}} f(t)\, e^{-i2\pi\eta t} dt$ is the Fourier transform of f. Certain examples and basic properties of the S-transform can be found in [7, 8, 9, 10, 11, 12, 13, 14].

1.1. **The fractional Fourier transform.** The fractional Fourier transform was introduced by Almeida [1] as a generalization of classical Fourier transform. It has several known applications in the signal processing and many other scientific fields. The a^{th} order FRFT (fractional Fourier transform) of a signal f is defined as

$$F_f^a(\xi) = \int_{\mathbb{R}} f(t)\, K_a(t,\xi)\, dt, \tag{1.6}$$

where the transform kernel is given by

$$K_a(t,\xi) = \begin{cases} A_\theta\, e^{i\pi(\xi^2 \cot\theta - 2\xi t \csc\theta + t^2 \cot\theta)}, & \text{if } \theta \neq j\pi \\ \delta(t-\xi), & \text{if } \theta = 2j\pi \\ \delta(t+\xi), & \text{if } \theta + \pi = 2j\pi, \end{cases} \tag{1.7}$$

where $A_\theta = \sqrt{1 - i\cot\theta}, \theta = a\frac{\pi}{2}$, $a \in [0,4)$, i is the complex unit, j is an integer and ξ is the fractional Fourier frequency. The inverse FRFT of equation (1.6) is

$$f(t) = \int_{\mathbb{R}} F_f^a(\xi)\overline{K_a(t,\xi)} d\xi.$$

1.2. **The fractional S-transform.** The fractional S-transform is used by Xu et.al.[19] in 2012 as a generalization of the S-transform. The a^{th} order continuous fractional S-transform (FRST) of a signal $f(t)$ is defined as

$$FRST_f^a(\tau,\xi) = \int_{\mathbb{R}} f(t) g(\tau - t, \xi) K_a(t,\xi) dt, \tag{1.8}$$

where the window function g is given by

$$g(t,\xi) = \frac{|\xi \csc\theta|^p}{k\sqrt{2\pi}} e^{-t^2(\xi\csc\theta)^{2p}/2k^2} \;;\; k,p > 0, \tag{1.9}$$

which satisfies the condition:

$$\int_{\mathbb{R}} g(t,\xi) dt = 1, \text{ for all } \xi \in \mathbb{R}_0. \tag{1.10}$$

Inverse fractional S-transform is defined as

$$f(t) = \int_{\mathbb{R}} \left[\int_{\mathbb{R}} FRST_f^a(\tau,\xi)\, d\tau \right] \overline{K_a(t,\xi)} d\xi.$$

The fractional S-transform depends on a parameter θ and can be interpreted as a rotation by an angle θ in the time-frequency plane. The parameter p and k can be used to adjust the window function space. The fractional S-transform has been studied on distribution spaces by Singh [9, 10, 11].

2. The fractional S-transform on BMO and Hardy spaces

The bounded mean oscillation space $BMO(\mathbb{R})$ also known as John-Nirenberg space was first introduced by F. John and L. Nirenberg in 1961[5]. It is the dual space of the real Hardy space H^1 and serves in many ways as a substitute space for L^∞. The $BMO(\mathbb{R})$ spaces play an important role in various areas of mathematics, for example many operators which are bounded on L^p, $1 < p < \infty$, but not on L^∞, are bounded when considered as operators on BMO.

Now, we recall the definitions of the BMO and Hardy spaces.

Definition 2.1. The bounded mean oscillation space $BMO(\mathbb{R})$ is defined in [16, p. 140] as the space of all Lebesgue integrable (locally) functions defined on $\mathbb{R}$ such that

$$\| f \|_{BMO} = \sup_{I \subset \mathbb{R}} \frac{1}{|I|} \int_I |f(x) - f_I| \, dx < \infty, \tag{2.1}$$

here the supremum is taken over all intervals I in $\mathbb{R}$ of measure $|I|$ and f_I stands for the mean of f on I, namely

$$f_I := \frac{1}{|I|} \int_I f(x) \, dx \leq \frac{1}{|I|} \int_I |f(x)| \, dx \leq m < \infty. \tag{2.2}$$

Definition 2.2. The Hardy space is defined in [16, pp. 90-91] as the space of all functions $f \in L^1(\mathbb{R})$ such that

$$\| f \|_{H^1} = \int_{\mathbb{R}} \sup_{t>0} |(f * \phi_t)(x)| \, dx < \infty, \tag{2.3}$$

where ϕ is any test function with $\int_{\mathbb{R}} \phi(x) dx \neq 0$ and $\phi_t(x) = t^{-1}\phi(x/t)$; $t > 0$, $x \in \mathbb{R}$.

Our main results in this section are as follows.

Lemma 2.1. *If $K_a(x, \xi)$ is the transform kernel defined in (1.7) and $\theta \neq j\pi$, then for any Lebesgue integrable (locally) function defined on $\mathbb{R}$ we have*

$$\|f(\cdot)K_a(\cdot, \xi)\|_{BMO} \leq |A_\theta| \left(\|f\|_{BMO} + 2m \right)$$

where m is a constant given in equation (2.2).

Proof. By using the inequality $|K_a(x,\xi)| \leq |A_\theta|$, we have

$$\begin{aligned}
&\|f(\cdot)K_a(\cdot,\xi)\|_{BMO}\\
&= \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left| f(x)K_a(x,\xi) - \frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt \right| dx\\
&= \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left| f(x)K_a(x,\xi) - \frac{K_a(x,\xi)}{|I|}\int_I f(t)dt + \frac{K_a(x,\xi)}{|I|}\int_I f(t)dt \right.\\
&\quad \left. - \frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt \right| dx\\
&\leq \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left| K_a(x,\xi)\left(f(x) - \frac{1}{|I|}\int_I f(t)dt \right) \right| dx\\
&\quad + \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left| \frac{K_a(x,\xi)}{|I|}\int_I f(t)dt \right| dx + \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left| \frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt \right| dx\\
&\leq |A_\theta|\,\|f\|_{BMO} + |A_\theta| \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I |f|_I\,dx + |A_\theta| \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \frac{1}{|I|}\int_I |f(t)|dtdx\\
&\leq |A_\theta| \left(\|f\|_{BMO} + \frac{1}{|I|}m|I| + \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I m\,dx \right)\\
&\leq |A_\theta| \left(\|f\|_{BMO} + \frac{1}{|I|}m|I| + \frac{1}{|I|}m|I| \right)\\
&= |A_\theta| \left(\|f\|_{BMO} + 2m \right).
\end{aligned}$$

□

Theorem 2.2. *For any fixed $\xi \in \mathbb{R}_0$ and $\theta \neq j\pi$, the operator $FRST_f^a : BMO(\mathbb{R}) \to BMO(\mathbb{R})$ is continuous and*

$$\| (FRST_f^a)(\cdot,\xi) \|_{BMO} \leq |A_\theta| \left(\|f\|_{BMO} + 2m \right).$$

Proof. Since

$$\begin{aligned}
&\left| (FRST_f^a)(\tau,\xi) - (FRST_f^a)_I(\tau,\xi) \right|\\
&= \left| \int_{\mathbb{R}} f(\tau-x)g(x,\xi)K_a(\tau-x,\xi)dx \right.\\
&\quad \left. - \frac{1}{|I|}\int_I\int_{\mathbb{R}} f(\alpha-x)g(x,\xi)K_a(\alpha-x,\xi)dx\,d\alpha \right|\\
&= \left| \int_{\mathbb{R}} f(\tau-x)g(x,\xi)K_a(\tau-x,\xi)dx \right.\\
&\quad \left. - \int_{\mathbb{R}} g(x,\xi)\left(\frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha \right) dx \right|\\
&= \left| \int_{\mathbb{R}} g(x,\xi)\left(f(\tau-x)K_a(\tau-x,\xi) - \frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha \right) dx \right|\\
&\leq \int_{\mathbb{R}} |g(x,\xi)| \left| f(\tau-x)K_a(\tau-x,\xi) - \frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha \right| dx.
\end{aligned}$$

Therefore,

$$
\begin{aligned}
&\| (FRST_f^a)(\cdot,\xi) \|_{BMO} \\
&= \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \left|(FRST_f^a)(\tau,\xi) - (FRST_f^a)_I(\tau,\xi)\right| d\tau \\
&\leq \sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \int_{\mathbb{R}} |g(x,\xi)| \Bigg| f(\tau-x)K_a(\tau-x,\xi) \\
&- \frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha\Bigg| dx\, d\tau \\
&= \int_{\mathbb{R}} |g(x,\xi)| \Bigg(\sup_{I\subset\mathbb{R}} \frac{1}{|I|}\int_I \Bigg| f(\tau-x)K_a(\tau-x,\xi) \\
&- \frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha\Bigg| d\tau\Bigg) dx \\
&= \int_{\mathbb{R}} |g(x,\xi)| \left(\sup_{J\subset\mathbb{R}} \frac{1}{|J|}\int_J \left| f(t)K_a(t,\xi) - \frac{1}{|J|}\int_J f(y)K_a(y,\xi)dy\right| dt\right) dx \\
&\qquad (\text{here } J = I - x \text{ for } x\in\mathbb{R}) \\
&= \|f(\cdot)K_a(\cdot,\xi)\|_{BMO} \int_{\mathbb{R}} |g(x,\xi)|dx \\
&= \|f(\cdot)K_a(\cdot,\xi)\|_{BMO}\,.
\end{aligned}
$$

Now by using above lemma we get

$$
\| (FRST_f^a)(\cdot,\xi) \|_{BMO} \quad \leq \quad |A_\theta| \left(\|f\|_{BMO} + 2m\right).
$$

□

Theorem 2.3. *Let $f \in L^1(\mathbb{R})$ such that*

$$
\sup_{t>0} \left| \int_{\mathbb{R}} f(x-y)\phi_t(y)dy \right| = \sup_{t>0} \int_{\mathbb{R}} |f(x-y)\phi_t(y)|\, dy < \infty. \tag{2.4}
$$

Then for any fixed $\xi \in \mathbb{R}_0$, the operator $FRST_f^a : H^1(\mathbb{R}) \to H^1(\mathbb{R})$ is continuous and

$$
\| (FRST)(\cdot,\xi) \|_{H^1} \leq |A_\theta| \parallel f \parallel_{H^1}\,.
$$

Proof. We have

$$
\begin{aligned}
&\| FRST_f^a(\cdot,\xi) \|_{H^1} \\
&= \int_{\mathbb{R}} \sup_{t>0} \left| \left(FRST_f^a(\cdot,\xi) * \phi_t(\cdot) \right)(\tau) \right| d\tau \\
&= \int_{\mathbb{R}} \sup_{t>0} \left| \left(\left(\int_{\mathbb{R}} f(\cdot - x) g(x,\xi) K_a(\cdot - x, \xi) dx \right) * \phi_t(\cdot) \right)(\tau) \right| d\tau \\
&= \int_{\mathbb{R}} \sup_{t>0} \left| \int_{\mathbb{R}} g(x,\xi) \left(\int_{\mathbb{R}} f(\tau - x - y) K_a(\tau - x - y, \xi) \phi_t(y) dy \right) dx \right| d\tau \\
&\leq \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(\tau - x - y) K_a(\tau - x - y, \xi) \phi_t(y)| \, dy d\tau \right) dx \\
&= \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(z) K_a(z,\xi) \phi_t(\tau - x - z)| \, dz d\tau \right) dx \\
&\leq |A_\theta| \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(z) \phi_t(\tau - x - z)| \, dz d\tau \right) dx \\
&= |A_\theta| \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} |(f * \phi_t)(\tau - x)| \, d\tau \right) dx \\
&\qquad \text{(by using equation (2.4) and then definition of convolution)} \\
&= |A_\theta| \, \|f\|_{H^1} \int_{\mathbb{R}} |g(x,\xi)| \, dx \\
&= |A_\theta| \, \|f\|_{H^1} .
\end{aligned}
$$

□

3. The fractional S-transform on weighted function spaces

Let us recall the definitions of relevant weighted function spaces.

Definition 3.1. A positive function κ defined on $\mathbb{R}$ is called a tempered weight function if there exist positive constants C and N such that

$$\kappa(\xi + \eta) \leq (1 + C|\xi|)^N \kappa(\eta) \text{ for all } \xi, \eta \in \mathbb{R}, \tag{3.1}$$

and the set of all such functions κ is denoted by $\mathscr{K}$. Certain examples and basic properties of weight function κ can be found in [4].

Definition 3.2. For $1 \leq p < \infty$, the weighted Lebesgue space $L^p_\kappa(\mathbb{R})$ is defined as the space of all measurable functions f on $\mathbb{R}$ such that

$$\| f \|_{L^p_\kappa} = \left(\int_{\mathbb{R}} |f(x)|^p \, \kappa(x) dx \right)^{1/p} < \infty.$$

Definition 3.3. The weighted bounded mean oscillation space $\mathrm{BMO}_\kappa(\mathbb{R})$ is defined as the space of all weighted Lebesgue integrable (locally) functions defined on $\mathbb{R}$ such that

$$\| f \|_{BMO_\kappa} = \sup_{I \subset \mathbb{R}} \frac{1}{|I|_\kappa} \int_I |f(x) - f_I| \, \kappa(x) dx < \infty,$$

where the supremum is taken over all intervals I in $\mathbb{R}$ and $|I|_\kappa = \int_I \kappa(x) dx$.

Definition 3.4. The weighted Hardy space is defined as the space of all functions $f \in L^1_\kappa(\mathbb{R})$ such that

$$\| f \|_{H^1_\kappa} = \int_{\mathbb{R}} \sup_{t>0} |(f * \phi_t)(x)| \, \kappa(x) dx < \infty.$$

where ϕ is any test function with $\int_{\mathbb{R}} \phi(x)dx \neq 0$ and $\phi_t(x) = t^{-1}\phi(x/t)$; $t > 0$, $x \in \mathbb{R}$.

Our main results on weighted function spaces are as follows.

Lemma 3.1. *If $g(x, \xi)$ is defined by (1.9) and $\theta \neq j\pi$, then for any fixed $\xi \in \mathbb{R}_0$ we have the following estimate*

$$\int_{\mathbb{R}} |g(x, \xi)|(1 + C|x|)^N dx \leq A_{\xi,N},$$

where N is positive constant, and $A_{\xi,N}$ is a constant depend on ξ and N.

Proof. Using the well-known inequality, $|x + y|^n \leq 2^{n-1}(|x|^n + |y|^n)$, $\forall\, x, y \in \mathbb{R}$, $\forall\, n \in \mathbb{N}$, we have

$$\begin{aligned}
&\int_{\mathbb{R}} |g(x, \xi)|(1 + C|x|)^N dx \\
&\leq \int_{\mathbb{R}} |g(x, \xi)| \, 2^{[N]} \left(1 + |Cx|^{[N]+1}\right) dx \\
&= 2^{[N]} \left(1 + C^{[N]+1} \int_{\mathbb{R}} |g(x, \xi)||x|^{[N]+1} dx\right) \\
&= 2^{[N]} \left(1 + 2\, C^{[N]+1} \int_0^\infty x^{[N]+1} g(x, \xi) dx\right) \\
&= 2^{[N]} \left(1 + 2\, C^{[N]+1} \frac{2^{([N]+1)/2} k^{[N]+1}}{2\sqrt{\pi}\, |\xi \csc\theta|^{p([N]+1)}} \Gamma([N]/2 + 1)\right) \\
&\leq A_{\xi,N}.
\end{aligned}$$

□

Lemma 3.2. *If $K_a(x, \xi)$ is the transform kernel defined in (1.7) and $\theta \neq j\pi$, then for any locally Lebesgue integrable function defined on $\mathbb{R}$ we have*

$$\|f(\cdot)K_a(\cdot, \xi)\|_{BMO_\kappa} \leq |A_\theta| \left(\|f\|_{BMO_\kappa} + 2m\right)$$

where m is a constant given in equation (2.2).

Proof. We have

$$
\begin{aligned}
&\|f(\cdot)K_a(\cdot,\xi)\|_{BMO_\kappa}\\
&=\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|f(x)K_a(x,\xi)-\frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt\right|\kappa(x)dx\\
&=\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|f(x)K_a(x,\xi)-\frac{K_a(x,\xi)}{|I|}\int_I f(t)dt+\frac{K_a(x,\xi)}{|I|}\int_I f(t)dt\right.\\
&\left.-\frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt\right|\kappa(x)dx\\
&=\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|K_a(x,\xi)\left(f(x)-\frac{1}{|I|}\int_I f(t)dt\right)+\frac{K_a(x,\xi)}{|I|}\int_I f(t)dt\right.\\
&\left.-\frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt\right|\kappa(x)dx\\
&\leq\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|K_a(x,\xi)\left(f(x)-\frac{1}{|I|}\int_I f(t)dt\right)\right|\kappa(x)dx\\
&+\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|\frac{K_a(x,\xi)}{|I|}\int_I f(t)dt\right|\kappa(x)dx\\
&+\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left|\frac{1}{|I|}\int_I f(t)K_a(t,\xi)dt\right|\kappa(x)\,dx\\
&\leq|A_\theta|\,\|f\|_{BMO_\kappa}+|A_\theta|\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I|f|_I\,\kappa(x)dx\\
&+|A_\theta|\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I\left(\frac{1}{|I|}\int_I|f(t)|dt\right)\kappa(x)dx\\
&\leq|A_\theta|\left(\|f\|_{BMO_\kappa}+\frac{1}{|I|_\kappa}m|I|_\kappa+\sup_{I\subset\mathbb{R}}\frac{1}{|I|_\kappa}\int_I m\,\kappa(x)\,dx\right)\\
&\leq|A_\theta|\left(\|f\|_{BMO_\kappa}+\frac{1}{|I|_\kappa}m|I|_\kappa+\frac{1}{|I|_\kappa}m|I|_\kappa\right)\\
&=|A_\theta|\left(\|f\|_{BMO_\kappa}+2m\right).
\end{aligned}
$$

□

Theorem 3.3. *For any fixed $\xi\in\mathbb{R}_0$, the operator $FRST_f^a : BMO_\kappa(\mathbb{R})\to BMO_\kappa(\mathbb{R})$ is continuous. Furthermore, we have*

$$\|\,(FRST_f^a)(\cdot,\xi)\,\|_{BMO_\kappa}\leq A_{\xi,N}\,|A_\theta|\left(\|f\|_{BMO_\kappa}+2m\right)$$

where m is a constant given in equation (2.2).

Proof.

$$
\begin{aligned}
&\| (FRST_f^a)(\cdot,\xi) \|_{BMO_\kappa} \\
&= \sup_{I\subset\mathbb{R}} \frac{1}{|I|_\kappa} \int_I \left|(FRST_f^a)(\tau,\xi) - (FRST_f^a)_I(\tau,\xi)\right| \kappa(\tau)d\tau \\
&= \sup_{I\subset\mathbb{R}} \frac{1}{|I|_\kappa} \int_I \Bigg| \int_{\mathbb{R}} g(x,\xi)\Bigg(f(\tau-x)K_a(\tau-x,\xi) \\
&- \frac{1}{|I|}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha\Bigg)dx\Bigg|\kappa(\tau)d\tau \\
&\le \int_{\mathbb{R}} |g(x,\xi)|\Bigg(\sup_{I\subset\mathbb{R}} \frac{1}{|I|_\kappa} \int_I \Bigg| f(\tau-x)K_a(\tau-x,\xi) \\
&- \frac{1}{|I|_\kappa}\int_I f(\alpha-x)K_a(\alpha-x,\xi)d\alpha\Bigg| k(\tau)d\tau\Bigg)dx \\
&= \int_{\mathbb{R}} |g(x,\xi)|\Bigg(\sup_{J\subset\mathbb{R}} \frac{1}{|J|_\kappa} \int_J \Bigg| f(y)K_a(y,\xi) - \frac{1}{|J|_\kappa}\int_J f(t)K_a(t,\xi)dt\Bigg|\kappa(x+y)dy\Bigg)dx \\
&\qquad (\text{here } J = I - x \text{ for } x\in\mathbb{R}) \\
&\le \int_{\mathbb{R}} |g(x,\xi)|\Bigg(\sup_{J\subset\mathbb{R}} \frac{1}{|J|_\kappa} \int_J \Bigg| f(y)K_a(y,\xi) \\
&- \frac{1}{|J|_\kappa}\int_J f(t)K_a(t,\xi)dt\Bigg|(1+C|x|)^N\kappa(y)dy\Bigg)dx \\
&= \|f(\cdot)K_a(\cdot,\xi)\|_{BMO_\kappa} \int_{\mathbb{R}} |g(x,\xi)|(1+C|x|)^N dx.
\end{aligned}
$$

So by using lemma 3.1 and 3.2, we get

$$
\| (FRST_f^a)(\cdot,\xi) \|_{BMO_\kappa} \le A_{\xi,N}\,|A_\theta|\left(\|f\|_{BMO_\kappa} + 2m\right).
$$

□

Theorem 3.4. *Let $f \in L^1(\mathbb{R})$ and satisfies the condition (2.4). Then, for any fixed $\xi \in \mathbb{R}_0$, the operator $FRST_f^a : H_\kappa^1(\mathbb{R}) \to H_\kappa^1(\mathbb{R})$ is continuous. Furthermore, we have*

$$
\| FRST_f^a(\cdot,\xi) \|_{H_\kappa^1} \le A_{\xi,N}\,|A_\theta|\,\|f\|_{H_\kappa^1}.
$$

Proof. We have

$$
\begin{aligned}
&\| FRST_f^a(\cdot,\xi) \|_{H_\kappa^1} \\
&= \int_{\mathbb{R}} \sup_{t>0} \left| \left(FRST_f^a(\cdot,\xi) * \phi_t(\cdot) \right)(\tau) \right| \kappa(\tau) d\tau \\
&= \int_{\mathbb{R}} \sup_{t>0} \left| \int_{\mathbb{R}} g(x,\xi) \left(\int_{\mathbb{R}} f(\tau - x - y) K_a(\tau - x - y, \xi) \phi_t(y) dy \right) dx \right| \kappa(\tau) d\tau \\
&\leq \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(\tau - x - y) K_a(\tau - x - y, \xi) \phi_t(y)| \, \kappa(\tau) dy \, d\tau \right) dx \\
&= \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(z - y) K_a(z - y, \xi) \phi_t(y)| \, \kappa(z + x) dz \, dy \right) dx \\
&\leq |A_\theta| \int_{\mathbb{R}} |g(x,\xi)| \left(\int_{\mathbb{R}} \sup_{t>0} \int_{\mathbb{R}} |f(z - y) \phi_t(y)| \, (1 + C|x|)^N \kappa(z) dz \right) dx \\
&= |A_\theta| \int_{\mathbb{R}} |g(x,\xi)| \, (1 + C|x|)^N \left(\int_{\mathbb{R}} \sup_{t>0} |(f * \phi_t)(z)| \, \kappa(z) dz \right) dx \\
&\qquad \text{(by using equation (2.4) and then definition of convolution)} \\
&= |A_\theta| \, \|f\|_{H_\kappa^1} \int_{\mathbb{R}} |g(x,\xi)| \, (1 + C|x|)^N dx.
\end{aligned}
$$

Now by using lemma 3.1 we get

$$\| FRST_f^a(\cdot,\xi) \|_{H_\kappa^1} \leq A_{\xi,N} \, |A_\theta| \, \|f\|_{H_\kappa^1}.$$

□

References

[1] L. B. Almeida : The fractional Fourier transform and time-frequency representations. *IEEE Trans. Signal Process.* **42(11)** (1994), 3084-3091.

[2] N. M. Chuong, D. V. Duong : Boundedness of the wavelet integral operator on weighted function spaces. *Russ. J. Math. Phys.* **20(3)** (2013), 268-275.

[3] C. Fefferman, E. M. Stein : H^p spaces of several variables, *Acta Math.* **129(1)** (1972), 137-193.

[4] L. Hörmander : *The Analysis of Linear Partial Differential Operators II.* Springer, Berlin (1983).

[5] F. John, L. Nirenberg : On functions of bounded mean oscillation . *Commun. Pure Appl. Math.* **14** (1961), 415-426.

[6] R. S. Pathak, S. K. Singh : Boundedness of the wavelet transform in certain function spaces. *J. Inequal. Pure Appl. Math.* **8(1)** (2007), Article 23

[7] S. K. Singh : The S-transform on spaces of type S. *Integral Transforms Spec. Funct.* **23(7)** (2012), 481-494.

[8] S. K. Singh : The S-transform on spaces of type W. *Integral Transforms Spec. Funct.* **23(12)** (2012), 891-899.

[9] S. K. Singh : The fractional S-transform of tempered ultradistibutions. *Investig. Math. Sci.* **2(2)** (2012), 315-325.

[10] S. K. Singh : The fractional S-transform on spaces of type S. *J. Math.* 2013. doi: 10.1155/2013/105848. Article ID 105848.

[11] S. K. Singh : The fractional S-transform on spaces of type W. *J. Pseudo-Differ. Oper. Appl.* **4(2)** (2013), 251-265.

[12] S. K. Singh : A new integral transform: Theory part. *Investig. Math. Sci.* **3(1)** (2013), 135-139.

[13] S. K. Singh : The S-Transform of distributions. *Sci. World J.* 2014. doi: 10.1155/2014/623294. Article ID 623294.

[14] S. K. Singh : Besov norms in terms of the S-transform. *Afr. Mat.* (2015). doi: 10.1007/s13370-015-0365-0.
[15] S. K. Singh, B. Kalita : The S-transform on Hardy spaces and its duals. *International J. Anal. Appl.* **7(2)** (2015), 171-178.
[16] E. M. Stein : *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton University Press, Princeton (1993).
[17] R. G. Stockwell, L. Mansinha, R. P. Lowe : Localization of the complex spectrum: The S transform. *IEEE Trans. Signal Process.* **44(4)** (1996), 998-1001.
[18] S. Ventosa, C. Simon, M. Schimmel, J. Dañobeitia, A. Mànuel : The S-transform from a wavelet point of view. *IEEE Trans. Signal Process.* **56(07)** (2008), 2771-2780.
[19] D. P. Xu, K. Guo : Fractional S transform -Part 1:Theory. *Applied Geophysics.* **9(1)** (2012), 73-79.

Department of Mathematics, Rajiv Gandhi University
Doimukh-791112, Arunachal Pradesh, India
[1] Email: kalitababy478@gmail.com
[2] Email: sks_math@yahoo.com
* Corresponding author